



## Exelixis Announces Initiation of the STELLAR-303 Phase 3 Pivotal Trial Evaluating XL092 in Patients with Metastatic Colorectal Cancer

June 21, 2022

*– STELLAR-303 is the first phase 3 pivotal trial evaluating XL092, a next-generation oral tyrosine kinase inhibitor –*

ALAMEDA, Calif.--(BUSINESS WIRE)--Jun. 21, 2022-- [Exelixis, Inc.](#) (Nasdaq: EXEL) today announced the initiation of STELLAR-303, a phase 3 pivotal trial evaluating XL092 in combination with atezolizumab versus regorafenib in patients with metastatic colorectal cancer (CRC) that is not microsatellite instability-high or mismatch repair-deficient, who have progressed after or are intolerant to the standard of care therapy. XL092 is a next-generation tyrosine kinase inhibitor (TKI) in development for multiple advanced tumor types.

“There is a significant need for new treatment options for the majority of metastatic CRC patients, who do not have microsatellite instability-high or mismatch-repair deficient disease and whose tumors do not respond to immunotherapy alone,” said Vicki L. Goodman, M.D., Executive Vice President, Product Development & Medical Affairs, and Chief Medical Officer, Exelixis. “Following recent promising data evaluating cabozantinib in combination with immunotherapies in colorectal cancer, we are thrilled to initiate our first phase 3 pivotal trial for XL092, our next-generation tyrosine kinase inhibitor. We look forward to learning more about how XL092 in combination with atezolizumab may benefit patients with metastatic colorectal cancer.”

STELLAR-303 is a global, multicenter, randomized phase 3 open-label study that will enroll approximately 600 patients with documented RAS status. Patients will be randomized 1:1 to receive either XL092 in combination with atezolizumab or regorafenib. The primary objective of the study is to evaluate the efficacy of the combination in patients with RAS wild-type disease; exploratory endpoints include examining efficacy in those with RAS-mutated disease. The primary endpoint is overall survival. Secondary endpoints include progression-free survival, objective response rate and duration of response per Response Evaluation Criteria in Solid Tumors version 1.1 as assessed by the investigator.

Previously [announced](#) results from two studies of cabozantinib in combination with immunotherapies for the treatment of advanced CRC supported Exelixis' decision to pursue clinical development of XL092 in this setting. The trial is sponsored by Exelixis, and Roche is supplying atezolizumab.

### About XL092

XL092 is a next-generation oral TKI that inhibits the activity of receptor tyrosine kinases implicated in cancer growth and spread, including VEGF receptors, MET, AXL and MER. These receptor tyrosine kinases are involved in both normal cellular function and in pathologic processes such as oncogenesis, metastasis, tumor angiogenesis and resistance to multiple therapies, including immune checkpoint inhibitors. In designing XL092, Exelixis sought to build upon its extensive experience with and the target profile of cabozantinib, the company's flagship medicine, while improving key characteristics, including pharmacokinetic half-life. XL092 is currently being developed for the treatment of advanced solid tumors, including genitourinary cancers, as a monotherapy and in combination with immune checkpoint inhibitors. XL092 is the first internally discovered Exelixis compound to enter the clinic following the company's reinitiation of drug-discovery activities.

### About Colorectal Cancer

Colorectal cancer is the third most common cancer and the third-leading cause of cancer-related deaths in the U.S. According to the American Cancer Society, about 150,000 new cases will be diagnosed and 53,000 people will die from the disease in 2022.<sup>1</sup> Colorectal cancer is most frequently diagnosed among people aged 65-74 and is more common in men and those of African American descent. Nearly a quarter of colorectal cancer cases are diagnosed at the metastatic stage, at which point the five-year survival rate is just 15%.<sup>2</sup> It has been estimated that approximately 40% of metastatic colorectal cancer cases exhibit a RAS mutation.<sup>3</sup>

### About CABOMETYX® (cabozantinib)

In the U.S., CABOMETYX tablets are approved for the treatment of patients with advanced renal cell carcinoma (RCC); for the treatment of patients with hepatocellular carcinoma (HCC) who have been previously treated with sorafenib; for patients with advanced RCC as a first-line treatment in combination with nivolumab; and for adult and pediatric patients 12 years of age and older with locally advanced or metastatic differentiated thyroid cancer (DTC) that has progressed following prior VEGFR-targeted therapy and who are radioactive iodine-refractory or ineligible. CABOMETYX tablets have also received regulatory approvals in the European Union and additional countries and regions worldwide. In 2016, Exelixis granted Ipsen exclusive rights for the commercialization and further clinical development of cabozantinib outside of the U.S. and Japan. In 2017, Exelixis granted exclusive rights to Takeda for the commercialization and further clinical development of cabozantinib for all future indications in Japan. Exelixis holds the exclusive rights to develop and commercialize cabozantinib in the U.S.

CABOMETYX is not indicated as a treatment for metastatic colorectal cancer CRC that is not microsatellite instability-high or mismatch repair-deficient.

### IMPORTANT SAFETY INFORMATION

#### WARNINGS AND PRECAUTIONS

**Hemorrhage:** Severe and fatal hemorrhages occurred with CABOMETYX. The incidence of Grade 3 to 5 hemorrhagic events was 5% in CABOMETYX patients in RCC, HCC, and DTC studies. Discontinue CABOMETYX for Grade 3 or 4 hemorrhage and prior to surgery as recommended. Do not administer CABOMETYX to patients who have a recent history of hemorrhage, including hemoptysis, hematemesis, or melena.

**Perforations and Fistulas:** Fistulas, including fatal cases, occurred in 1% of CABOMETYX patients. Gastrointestinal (GI) perforations, including fatal cases, occurred in 1% of CABOMETYX patients. Monitor patients for signs and symptoms of fistulas and perforations, including abscess and sepsis. Discontinue CABOMETYX in patients who experience a Grade 4 fistula or a GI perforation.

**Thrombotic Events:** CABOMETYX increased the risk of thrombotic events. Venous thromboembolism occurred in 7% (including 4% pulmonary embolism) and arterial thromboembolism in 2% of CABOMETYX patients. Fatal thrombotic events occurred in CABOMETYX patients. Discontinue CABOMETYX in patients who develop an acute myocardial infarction or serious arterial or venous thromboembolic events that require medical intervention.

**Hypertension and Hypertensive Crisis:** CABOMETYX can cause hypertension, including hypertensive crisis. Hypertension was reported in 37% (16% Grade 3 and <1% Grade 4) of CABOMETYX patients. Do not initiate CABOMETYX in patients with uncontrolled hypertension. Monitor blood pressure regularly during CABOMETYX treatment. Withhold CABOMETYX for hypertension that is not adequately controlled with medical management; when controlled, resume at a reduced dose. Permanently discontinue CABOMETYX for severe hypertension that cannot be controlled with anti-hypertensive therapy or for hypertensive crisis.

**Diarrhea:** Diarrhea occurred in 62% of CABOMETYX patients. Grade 3 diarrhea occurred in 10% of CABOMETYX patients. Monitor and manage patients using antidiarrheals as indicated. Withhold CABOMETYX until improvement to  $\leq$  Grade 1, resume at a reduced dose.

**Palmar-Plantar Erythrodysesthesia (PPE):** PPE occurred in 45% of CABOMETYX patients. Grade 3 PPE occurred in 13% of CABOMETYX patients. Withhold CABOMETYX until improvement to Grade 1 and resume at a reduced dose for intolerable Grade 2 PPE or Grade 3 PPE.

**Hepatotoxicity:** CABOMETYX in combination with nivolumab can cause hepatic toxicity with higher frequencies of Grades 3 and 4 ALT and AST elevations compared to CABOMETYX alone. Monitor liver enzymes before initiation of and periodically throughout treatment. Consider more frequent monitoring of liver enzymes than when the drugs are administered as single agents. For elevated liver enzymes, interrupt CABOMETYX and nivolumab and consider administering corticosteroids.

With the combination of CABOMETYX and nivolumab, Grades 3 and 4 increased ALT or AST were seen in 11% of patients. ALT or AST >3 times ULN (Grade  $\geq$ 2) was reported in 83 patients, of whom 23 (28%) received systemic corticosteroids; ALT or AST resolved to Grades 0-1 in 74 (89%). Among the 44 patients with Grade  $\geq$ 2 increased ALT or AST who were rechallenged with either CABOMETYX (n=9) or nivolumab (n=11) as a single agent or with both (n=24), recurrence of Grade  $\geq$ 2 increased ALT or AST was observed in 2 patients receiving CABOMETYX, 2 patients receiving nivolumab, and 7 patients receiving both CABOMETYX and nivolumab. Withhold and resume at a reduced dose based on severity.

**Adrenal Insufficiency:** CABOMETYX in combination with nivolumab can cause primary or secondary adrenal insufficiency. For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment, including hormone replacement as clinically indicated. Withhold CABOMETYX and/or nivolumab and resume CABOMETYX at a reduced dose depending on severity.

Adrenal insufficiency occurred in 4.7% (15/320) of patients with RCC who received CABOMETYX with nivolumab, including Grade 3 (2.2%), and Grade 2 (1.9%) adverse reactions. Adrenal insufficiency led to permanent discontinuation of CABOMETYX and nivolumab in 0.9% and withholding of CABOMETYX and nivolumab in 2.8% of patients with RCC.

Approximately 80% (12/15) of patients with adrenal insufficiency received hormone replacement therapy, including systemic corticosteroids. Adrenal insufficiency resolved in 27% (n=4) of the 15 patients. Of the 9 patients in whom CABOMETYX with nivolumab was withheld for adrenal insufficiency, 6 reinstated treatment after symptom improvement; of these, all (n=6) received hormone replacement therapy and 2 had recurrence of adrenal insufficiency.

**Proteinuria:** Proteinuria was observed in 8% of CABOMETYX patients. Monitor urine protein regularly during CABOMETYX treatment. For Grade 2 or 3 proteinuria, withhold CABOMETYX until improvement to  $\leq$  Grade 1 proteinuria, resume CABOMETYX at a reduced dose. Discontinue CABOMETYX in patients who develop nephrotic syndrome.

**Osteonecrosis of the Jaw (ONJ):** ONJ occurred in <1% of CABOMETYX patients. ONJ can manifest as jaw pain, osteomyelitis, osteitis, bone erosion, tooth or periodontal infection, toothache, gingival ulceration or erosion, persistent jaw pain, or slow healing of the mouth or jaw after dental surgery. Perform an oral examination prior to CABOMETYX initiation and periodically during treatment. Advise patients regarding good oral hygiene practices. Withhold CABOMETYX for at least 3 weeks prior to scheduled dental surgery or invasive dental procedures, if possible. Withhold CABOMETYX for development of ONJ until complete resolution, resume at a reduced dose.

**Impaired Wound Healing:** Wound complications occurred with CABOMETYX. Withhold CABOMETYX for at least 3 weeks prior to elective surgery. Do not administer CABOMETYX for at least 2 weeks after major surgery and until adequate wound healing. The safety of resumption of CABOMETYX after resolution of wound healing complications has not been established.

**Reversible Posterior Leukoencephalopathy Syndrome (RPLS):** RPLS, a syndrome of subcortical vasogenic edema diagnosed by characteristic findings on MRI, can occur with CABOMETYX. Evaluate for RPLS in patients presenting with seizures, headache, visual disturbances, confusion, or altered mental function. Discontinue CABOMETYX in patients who develop RPLS.

**Thyroid Dysfunction:** Thyroid dysfunction, primarily hypothyroidism, has been observed with CABOMETYX. Based on the safety population, thyroid dysfunction occurred in 19% of patients treated with CABOMETYX, including Grade 3 in 0.4% of patients.

Patients should be assessed for signs of thyroid dysfunction prior to the initiation of CABOMETYX and monitored for signs and symptoms of thyroid dysfunction during CABOMETYX treatment. Thyroid function testing and management of dysfunction should be performed as clinically indicated.

**Hypocalcemia:** CABOMETYX can cause hypocalcemia. Based on the safety population, hypocalcemia occurred in 13% of patients treated with CABOMETYX, including Grade 3 in 2% and Grade 4 in 1% of patients. Laboratory abnormality data were not collected in CABOSUN.

In COSMIC-311, hypocalcemia occurred in 36% of patients treated with CABOMETYX, including Grade 3 in 6% and Grade 4 in 3% of patients.

Monitor blood calcium levels and replace calcium as necessary during treatment. Withhold and resume at reduced dose upon recovery or permanently discontinue CABOMETYX depending on severity.

**Embryo-Fetal Toxicity:** CABOMETYX can cause fetal harm. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Verify the pregnancy status of females of reproductive potential prior to initiating CABOMETYX and advise them to use effective contraception during treatment and for 4 months after the last dose.

## ADVERSE REACTIONS

The most common ( $\geq 20\%$ ) adverse reactions are:

CABOMETYX as a single agent: diarrhea, fatigue, PPE, decreased appetite, hypertension, nausea, vomiting, weight decreased, constipation.

CABOMETYX in combination with nivolumab: diarrhea, fatigue, hepatotoxicity, PPE, stomatitis, rash, hypertension, hypothyroidism, musculoskeletal pain, decreased appetite, nausea, dysgeusia, abdominal pain, cough, and upper respiratory tract infection.

## DRUG INTERACTIONS

**Strong CYP3A4 Inhibitors:** If coadministration with strong CYP3A4 inhibitors cannot be avoided, reduce the CABOMETYX dosage. Avoid grapefruit or grapefruit juice.

**Strong CYP3A4 Inducers:** If coadministration with strong CYP3A4 inducers cannot be avoided, increase the CABOMETYX dosage. Avoid St. John's wort.

## USE IN SPECIFIC POPULATIONS

**Lactation:** Advise women not to breastfeed during CABOMETYX treatment and for 4 months after the final dose.

**Hepatic Impairment:** In patients with moderate hepatic impairment, reduce the CABOMETYX dosage. Avoid CABOMETYX in patients with severe hepatic impairment.

**Please see accompanying full Prescribing Information** <https://www.cabometryx.com/downloads/CABOMETYXUSPI.pdf>.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.FDA.gov/medwatch](http://www.FDA.gov/medwatch) or call 1-800-FDA-1088.

## About Exelixis

Founded in 1994, Exelixis, Inc. (Nasdaq: EXEL) is a commercially successful, oncology-focused biotechnology company that strives to accelerate the discovery, development and commercialization of new medicines for difficult-to-treat cancers. Following early work in model system genetics, we established a broad drug discovery and development platform that has served as the foundation for our continued efforts to bring new cancer therapies to patients in need. Our discovery efforts have resulted in four commercially available products, CABOMETYX<sup>®</sup> (cabozantinib), COMETRIQ<sup>®</sup> (cabozantinib), COTELLIC<sup>®</sup> (cobimetinib) and MINNEBRO<sup>®</sup> (esaxerenone), and we have entered into partnerships with leading pharmaceutical companies to bring these important medicines to patients worldwide. Supported by revenues from our marketed products and collaborations, we are committed to prudently reinvesting in our business to maximize the potential of our pipeline. We are supplementing our existing therapeutic assets with targeted business development activities and internal drug discovery – all to deliver the next generation of Exelixis medicines and help patients recover stronger and live longer. Exelixis is a member of the Standard & Poor's (S&P) MidCap 400 index, which measures the performance of profitable mid-sized companies. For more information about Exelixis, please visit [www.exelixis.com](http://www.exelixis.com), follow [@ExelixisInc](https://twitter.com/ExelixisInc) on Twitter or like [Exelixis, Inc.](https://www.facebook.com/ExelixisInc) on Facebook.

## Forward-looking Statements

This press release contains forward-looking statements, including, without limitation, statements related to: the clinical and therapeutic potential of XL092 in combination with atezolizumab as a treatment for patients with metastatic colorectal cancer; and Exelixis' plans to reinvest in its business to maximize the potential of the company's pipeline, including through targeted business development activities and internal drug discovery. Any statements that refer to expectations, projections or other characterizations of future events or circumstances are forward-looking statements and are based upon Exelixis' current plans, assumptions, beliefs, expectations, estimates and projections. Forward-looking statements involve risks and uncertainties. Actual results and the timing of events could differ materially from those anticipated in the forward-looking statements as a result of these risks and uncertainties, which include, without limitation: the potential failure of the combination of XL092 and atezolizumab to demonstrate safety and/or efficacy in STELLAR-303; uncertainties inherent in the product development process; complexities and the unpredictability of the regulatory review and approval processes in the U.S. and elsewhere; Exelixis' and Roche's continuing compliance with applicable legal and regulatory requirements; the continuing COVID-19 pandemic and other global events and their impact on Exelixis' research and development operations, including Exelixis' ability to initiate new clinical trials and clinical trial sites, enroll clinical trial patients, conduct trials per protocol, and conduct drug research and discovery operations and related activities; the costs of conducting clinical trials; Exelixis' dependence on third-party vendors for the development, manufacture and supply of XL092; Exelixis' ability to protect its intellectual property rights; market competition; changes in economic and business conditions; and other factors affecting Exelixis and its development programs discussed under the caption "Risk Factors" in Exelixis' Quarterly Report on Form 10-Q filed with the Securities and Exchange Commission (SEC) on May 10, 2022, and in Exelixis' future filings with the SEC. All forward-looking statements in this press release are based on information available to Exelixis as of the date of this press release, and Exelixis undertakes no obligation to update or revise any forward-looking statements contained herein, except as required by law.

*Exelixis, the Exelixis logo, CABOMETYX and COMETRIQ are registered U.S. trademarks of Exelixis.*

*COTELLIC is a registered trademark of Genentech, Inc.*

*MINNEBRO is a registered trademark of Daiichi Sankyo Company, Limited.*

<sup>1</sup> Cancer Facts and Figures 2022. American Cancer Society website. Available at: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2022/2022-cancer-facts-and-figures.pdf>. Accessed June 2022.

<sup>2</sup> Cancer Stat Facts: Colorectal Cancer. SEER website. Available at: <https://seer.cancer.gov/statfacts/html/colorect.html>. Accessed June 2022.

<sup>3</sup> RAS in Colorectal Cancer: ESMO Biomarker Factsheet. OncologyPRO website. Available at <https://oncologypro.esmo.org/education-library>

[/factsheets-on-biomarkers/ras-in-colorectal-cancer](#). Accessed June 2022.

View source version on [businesswire.com](https://www.businesswire.com/news/home/20220620005441/en/): <https://www.businesswire.com/news/home/20220620005441/en/>

**Investors:**

*Susan Hubbard*  
*EVP, Public Affairs and*  
*Investor Relations*  
*Exelixis, Inc.*  
*(650) 837-8194*  
[shubbard@exelixis.com](mailto:shubbard@exelixis.com)

**Media:**

*Lindsay Treadway*  
*Executive Director, Public Affairs*  
*and Advocacy Relations*  
*Exelixis, Inc.*  
*(650) 837-7522*  
[ltreadway@exelixis.com](mailto:ltreadway@exelixis.com)

Source: Exelixis, Inc.